

# ON THE ADDITIVITY OF LATTICE COMPLETENESS

to the memory of Maurice Audin

ISRAEL HALPERIN AND MARIA WONENBURGER

**1. Introduction.** It was shown in [1, Theorem 4.3] that upper  $\aleph$ -continuity<sup>1</sup> is additive in the following sense:

(1.1) *Suppose that  $[0, a]$ ,  $[0, b]$  are upper  $\aleph$ -continuous in a relatively complemented modular lattice. Then  $[0, a \cup b]$  is upper  $\aleph$ -continuous provided that  $[0, a \cup b]$  is upper  $\aleph$ -complete.*

But it may happen that  $[0, a]$ ,  $[0, b]$  are both upper  $\aleph$ -complete (both may even be von Neumann geometries with a perspective to  $b$ ) and yet  $[0, a \cup b]$  is *not* upper  $\aleph$ -complete. In fact there are von Neumann rings  $\mathcal{R}$  for which the lattice  $\bar{R}_{\mathcal{S}}$ , with  $\mathcal{S} = \mathcal{R}_2$ , is not even upper  $\aleph_0$ -complete (see the Remark preceding Definition 3.1)

With a modest supplementary condition however, additivity of upper  $\aleph$ -completeness does hold, as we show in this paper.

**2. Terminology and notation.** We shall use the notation of [1], [2], and [4].

I will denote a set of indices  $\alpha$  and  $\bar{I}$  will denote the cardinal power of  $I$ .

$\aleph$  will denote an infinite cardinal,  $\Omega$  will denote the least ordinal number whose corresponding cardinal power is  $\aleph$ .

A lattice is called *upper  $\aleph$ -complete* if the union  $a = \bigcup (a_\alpha | \alpha \in I)$  exists whenever  $\bar{I} \leq \aleph$ , and is called *upper  $\aleph$ -continuous* if for every  $b$ :  $b \cap a = \bigcup ((b \cap \bigcup (a_\alpha | \alpha \in F)) | \text{all finite } F \subset I)$ , with dual definitions for *lower  $\aleph$ -completeness* and *lower  $\aleph$ -continuity*. The lattice is called  *$\aleph$ -complete*, respectively  *$\aleph$ -continuous* if it is both upper and lower  $\aleph$ -continuous.

A complemented modular lattice  $L$  is called an  *$\aleph$ -von Neumann-geometry* if it is  $\aleph$ -complete and  $\aleph$ -continuous (irreducibility is *not* assumed).

If we omit the  $\aleph$  in any of these designations, this implies that the lattice  $L$  has the corresponding  $\aleph$ -property for all  $\aleph$ .

If  $\mathcal{R}$  is an associative regular ring (not necessarily with unit element) then  $\bar{R}_{\mathcal{R}}$  denotes the relatively complemented modular lattice of its principal right ideals, ordered by inclusion.  $\mathcal{R}$  is called an  *$\aleph$ -von Neumann-ring*, respectively a *von Neumann ring*, according as  $\bar{R}_{\mathcal{R}}$  is an  $\aleph$ -von

---

Received December 28, 1961. Dr. Wonenburger is a postdoctorate Fellow (of the National Research Council of Canada) at Queen's University.

<sup>1</sup> Terminology and notation are explained in section 2 below.

Neumann-geometry, respectively a von Neumann geometry.

In any relatively complemented modular lattice, if  $a \geq b$  then  $[a - b]$  will denote an arbitrary (but fixed) element such that  $[a - b] \dot{\cup} b = a$  (the dot indicates that the summands in the union are independent). We write  $a \sim b$  to denote:  $a$  is perspective to  $b$ , and  $a \lesssim b$  to denote:  $a \sim b_1$  for some  $b_1 \leq b$ . Elements  $a, b$  are called *completely disjoint*, (notation:  $(a, b)P$ ) if:  $a_1 \sim b_1, a_1 \leq a, b_1 \leq b$  together imply  $a_1 = 0$ .

**3. The additivity of completeness theorem.**

In this section  $a, b, c, \dots, x_\alpha, \dots$  will denote elements in a given relatively complemented modular lattice  $L$ .

If  $[0, a \cup c]$  is upper  $\aleph$ -complete we shall write  $u(a, c, \aleph)$  to mean:

(3.1) Whenever  $x_\alpha \leq a \cup c$  for all  $\alpha \in I$  (with  $\bar{I} \leq \aleph$ ) and

$$a \cap (\bigcup(x_\beta | \beta \in F)) = 0$$

for all finite  $F \subset I$ , then  $a \cap (\bigcup(x_\alpha | \alpha \in I)) = 0$ .

It is important to note: if  $u(a, c, \aleph)$  holds then  $u(a', c', \aleph)$  holds for all  $a' \leq a, c' \leq c$ .

Clearly, if  $[0, a \cup c]$  is upper  $\aleph$ -complete and upper  $\aleph$ -continuous then  $u(a, c, \aleph)$  does hold.

Similarly, if  $[0, a \cup c]$  is lower  $\aleph$ -complete we shall write  $l(a, c, \aleph)$  to denote:

(3.1) Whenever  $x_\alpha \leq a \cup c$  for all  $\alpha \in I$  (with  $\bar{I} \leq \aleph$ ) and

$$a \cup (\bigcap(x_\beta | \beta \in F)) = a \cup c$$

for all finite  $F \subset I$ , then  $a \cup (\bigcap(x_\alpha | \alpha \in I)) = a \cup c$ .

It is important to note: if  $l(a, c, \aleph)$  holds then  $l(a', c', \aleph)$  holds for all  $a' \leq a, c' \leq c$ .

Clearly, if  $[0, a \cup c]$  is lower  $\aleph$ -complete and lower  $\aleph$ -continuous then  $l(a, c, \aleph)$  does hold.

**LEMMA 3.1.** *Suppose that each of  $[0, a \cup b], [0, b \cup c], [0, a \cup c]$  is upper  $\aleph$ -complete and suppose that  $u(a, c, \aleph)$  holds. Then  $[0, a \cup b \cup c]$  is upper  $\aleph$ -complete.*

*Proof.* We may suppose that  $\{a, b, c\}$  is an independent set, for if  $c, b$  are replaced by  $[c - (a \cap c)]$  and  $[b - (b \cap (a \cup c))]$  respectively the hypotheses of Lemma 3.1 continue to hold and the conclusion is not changed.

Using transfinite induction, we may suppose that Lemma 3.1 holds

for all  $\aleph' < \aleph$ . We may therefore assume that  $x_\alpha$  is given,  $\leq a \cup b \cup c$  for all  $0 < \alpha < \Omega$ , that  $\mathbf{U}(x_\alpha | \alpha \leq \beta)$  exists for all  $\beta < \Omega$  and we need only show that  $\mathbf{U}(x_\alpha | \alpha < \Omega)$  exists.

We may suppose  $x_\alpha \leq x_\beta$  for  $\alpha \leq \beta < \Omega$  (by replacing the original  $x_\alpha$  by  $\mathbf{U}(x_\beta | \beta \leq \alpha)$  for all  $(\alpha < \Omega)$ ).

Set  $\bar{x}_0 = \mathbf{U}((x_\alpha \cap (a \cap b)) | \alpha < \Omega)$  (this union exists since, by hypothesis,  $[0, a \cup b]$  is upper  $\aleph$ -complete). Set  $\bar{x}_\alpha = \bar{x}_0 \cup x_\alpha$  for  $0 < \alpha < \Omega$  and observe that  $\bar{x}_\beta \leq \bar{x}_\alpha$  for all  $0 \leq \beta \leq \alpha < \Omega$ .

Set  $y_0 = \bar{x}_0$  and  $y_\alpha = [\bar{x}_\alpha - \mathbf{U}(\bar{x}_\beta | 0 \leq \beta < \alpha)]$  for  $0 < \alpha < \Omega$ . Then  $\mathbf{U}(y_\beta | 0 \leq \beta < \alpha) = \mathbf{U}(\bar{x}_\beta | 0 \leq \beta < \alpha)$  for all  $0 < \alpha < \Omega$ , as may be verified easily by transfinite induction.

Clearly, we need only show that  $\mathbf{U}(y_\alpha | 0 \leq \alpha < \Omega)$  exists. Hence it is sufficient to show that  $\mathbf{U}_\alpha y_\alpha$  exists, where (for the rest of this proof) we write  $\mathbf{U}_\alpha$  to mean  $\mathbf{U}_{0 < \alpha < \Omega}$  (note:  $0 \leq \alpha < \Omega$  has been replaced by  $0 < \alpha < \Omega$ ).

Set  $u = (a \cup (\mathbf{U}_\alpha((a \cup y_\alpha) \cap (b \cup c)))) \cap (b \cup (\mathbf{U}_\alpha((b \cup y_\alpha) \cap (a \cup c))))$  (this union exists since, by hypothesis,  $[0, b \cup c]$  and  $[0, a \cup c]$  are upper  $\aleph$ -complete). We observe that  $u \geq y_\beta$  for all  $0 < \beta < \Omega$  since each factor of  $u$  has this property: for fixed  $\beta$ ,  $a \cup (\mathbf{U}_\alpha((a \cup y_\alpha) \cap (b \cup c))) \geq a \cup ((a \cup y_\beta) \cap (b \cup c)) = (a \cup y_\beta) \cap (a \cup b \cup c) = a \cup y_\beta \geq y_\beta$ .

We shall show that  $u$  is the desired union  $\mathbf{U}_\alpha y_\alpha$ . It is clearly sufficient to show for every  $w$ : if  $u \geq w \geq y_\alpha$  for all  $0 < \alpha < \Omega$  then  $u \leq w$ .

Since  $a \cup y_\alpha \leq a \cup w$  and  $b \cup y_\alpha \leq b \cup w$  for all  $0 < \alpha < \Omega$ ,

$$\begin{aligned} u &\leq (a \cup ((a \cup w) \cap (b \cup c))) \cap (b \cup ((b \cup w) \cap (a \cup c))) \\ &= (a \cup w) \cap (b \cup w) = w \cup (a \cap (b \cup w)). \end{aligned}$$

It is therefore sufficient to show that  $a \cap (b \cup w) \leq w$ . We shall show that  $a \cap (b \cup u) = 0$ ; this will imply:

$$a \cap (b \cup w) \leq a \cap (b \cup u) = 0 \leq w.$$

$$\text{Now } b \cup u = (a \cup b \cup (\mathbf{U}_\alpha(a \cup y_\alpha) \cap (b \cup c))) \cap (b \cup (\mathbf{U}_\alpha((b \cup y_\alpha) \cap (a \cup c)))).$$

$$\begin{aligned} a \cap (b \cup u) &= a \cap (b \cup (\mathbf{U}_\alpha((b \cup y_\alpha) \cap (a \cup c)))) \\ &= a \cap ((b \cap (a \cup c)) \cup (\mathbf{U}_\alpha((b \cup y_\alpha) \cap (a \cup c)))) \\ &= a \cap (\mathbf{U}_\alpha((b \cup y_\alpha) \cap (a \cup c))). \end{aligned}$$

Since  $u(a, c, \aleph)$  is assumed to hold we need only show:

$$a \cap (\mathbf{U}(((b \cup y_\alpha) \cap (a \cup c)) | \alpha = \alpha_1, \dots, \alpha_m)) = 0$$

for every finite set of indices  $0 < \alpha_1 < \alpha_2 < \dots < \alpha_m < \Omega$ .

Hence it is sufficient to show that

$$a \cap (b \cup (\mathbf{U}(y_\alpha | \alpha = \alpha_1, \dots, \alpha_m))) = 0,$$

and so it is sufficient to show that

$$(3.2) \quad (a \cup b) \cap (\mathbf{U}(y_\alpha | \alpha = \alpha_1, \dots, \alpha_m)) = 0 .$$

For this purpose, we note:  $y_\alpha \cap (\mathbf{U}(y_\beta | 0 \leq \beta < \alpha) = 0$  for all  $0 < \alpha < \Omega$ . This implies that  $\{y_\alpha | \alpha = 0, \alpha_1, \dots, \alpha_m\}$  is an independent set and hence  $y_0 \cap (\mathbf{U}(y_\alpha | \alpha = \alpha_1, \dots, \alpha_m)) = 0$ . This implies (3.2) since the left side of (3.2) is  $\leq y_0$ . Thus Lemma 3.1 is proved.

**COROLLARY 1.** *Suppose that  $[0, a_i \cup a_j]$  is upper  $\aleph$ -complete for  $i, j = 1, \dots, m$  for some finite integer  $m$  and suppose that  $u(a_i, a_j, \aleph)$  holds whenever  $i < j$ . Then  $[0, a_1 \cup \dots \cup a_m]$  is upper  $\aleph$ -complete.*

*Proof.* If  $m \leq 2$  the conclusion is part of the hypotheses. Suppose that  $m > 2$  and that the Corollary is known to hold with  $m - 1$  in place of  $m$ ; then Lemma 3.1 can be applied (with  $a = a_1, b = a_3 \cup \dots \cup a_m$  and  $c = a_2$ ) to show that the Corollary holds for  $m$  itself. By induction on  $m$ , the Corollary is established.

**COROLLARY 2.** *Suppose that  $[0, a_i \cup a_j]$  is upper  $\aleph$ -complete and upper  $\aleph$ -continuous for  $i, j = 1, \dots, m$  for some finite integer  $m$ . Then  $[0, a_1 \cup \dots \cup a_m]$  is upper  $\aleph$ -complete and upper  $\aleph$ -continuous.*

*Proof.* Since upper  $\aleph$ -continuity of  $[0, a_i \cup a_j]$  implies that  $u(a_i, a_j, \aleph)$  holds, Corollary 1 shows that  $[0, a_1 \cup \dots \cup a_m]$  is upper  $\aleph$ -complete. The upper  $\aleph$ -continuity then follows from [1, Theorem 4.3].

**LEMMA 3.2.** *Suppose that  $a = a_1 \cup a_2 \cup \dots \cup a_m$  and  $a_i \leq a_1 \cup \dots \cup a_{i-1}$  for  $1 < i \leq m$ . Then  $a$  can be expressed in the form:*

$$(3.3) \quad a_1 \dot{\cup} \bar{a}_2 \dot{\cup} \dots \dot{\cup} \bar{a}_n \text{ for some } n \geq m \text{ and elements } \bar{a}_2, \dots, \bar{a}_n \text{ such that } \bar{a}_i \leq a_1 \text{ for all } 1 < i \leq n.$$

Moreover  $\bar{a}_2$  may be taken to coincide with  $a_2$  if  $a_1 \cap a_2 = 0$ .

*Proof.* Lemma 3.2 holds trivially if  $m = 1$  and also if  $m = 2$  and  $a_1 \cap a_2 = 0$ . We may therefore suppose (by induction) that  $m > 1$  and that  $b = a_1 \cup \dots \cup a_{m-1}$  has the form (3.3).

We can replace  $a_m$  by  $[a_m - (a_m \cap b)]$  since the hypotheses of Lemma 3.2 continue to hold and the conclusion is not changed. After this change,

$$a_m \cap b = a_m \cap (a_1 \dot{\cup} \bar{a}_2 \dot{\cup} \dots \dot{\cup} \bar{a}_n) = 0 .$$

Since  $a_m \leq a_1 \dot{\cup} \bar{a}_2 \dot{\cup} \dots \dot{\cup} \bar{a}_n$  there is a perspectivity mapping  $\varphi$  of  $[0, a_m]$  with  $\varphi(a_m) \leq b$ . Then

$$a_m = a_{m,1} \dot{\cup} a_{m,2} \dot{\cup} \dots \dot{\cup} a_{m,n}$$

where

$$\varphi(a_{m,1}) = \varphi(a_m) \cap a_1,$$

and for  $1 < i \leq n$ ,

$$\begin{aligned} \varphi(a_{m,i}) &= [(\varphi(a_m) \cap (a_1 \dot{\cup} \bar{a}_2 \dot{\cup} \dots \dot{\cup} \bar{a}_i)) \\ &\quad - (\varphi(a_m) \cap (a_1 \dot{\cup} \bar{a}_2 \dot{\cup} \dots \dot{\cup} \bar{a}_{i-1}))]. \end{aligned}$$

Obviously,  $a_{m,1} \lesssim a_1$ . If  $i > 1$  then  $a_{m,i} \sim \varphi(a_{m,i})$ ;  $\varphi(a_{m,i}) \lesssim \bar{a}_i$ ;  $\bar{a}_i \lesssim a_i$ ; and  $a_{m,i} \cap (\varphi(a_{m,i}) \cup \bar{a}_i \cup a_1) = 0$ ; these facts imply that  $a_{m,i} \lesssim a_1$  (use (2.2) of [1]). The conclusion of Lemma 3.2 now follows at once.

LEMMA 3.3. *Suppose that*

- (i)  $a = a_1 \cup a_2 \cup \dots \cup a_m$  for some finite  $m \geq 2$ ,
- (ii)  $a_2 \sim a_1$ ,
- (iii)  $a_i \lesssim a_1 \cup \dots \cup a_{i-1}$  for  $2 < i \leq m$ ,
- (iv)  $[0, a_1 \cup a_2]$  is upper  $\aleph$ -complete,
- (v)  $u(a_1, a_2, \aleph)$  holds.

Then  $[0, a]$  is upper  $\aleph$ -complete.

*Proof.* Applying Lemma 3.2, and using a new  $m$  and new elements  $a_3, \dots, a_m$  we may suppose that (i), (iii) hold in the strengthened form:  $a = a_1 \dot{\cup} a_2 \dot{\cup} \dots \dot{\cup} a_m$  and  $a_i \lesssim a_1$  for  $2 < i \leq m$ .

Suppose that  $1 \leq i < j \leq m$ . If  $i \neq 2$  then  $a_j \lesssim a_2$  (because of (ii)) and there is a perspectivity mapping  $\varphi$  of  $[0, a_i \cup a_j]$  with  $\varphi(a_i) \leq a_1$  and  $\varphi(a_j) \leq a_2$ . Hence  $[0, a_i \cup a_j]$  is upper  $\aleph$ -complete and  $u(a_i, a_j, \aleph)$  holds in this case.

If  $i = 2$  there is a perspectivity mapping  $\varphi$  of  $[0, a_2 \cup a_j]$  with  $\varphi(a_2) = a_1$ ,  $\varphi(a_j) = a_j$ ; the result for  $[0, a_1 \cup a_j]$  obtained previously now implies:  $[0, a_2 \cup a_j]$  is upper  $\aleph$ -complete and  $u(a_2, a_j, \aleph)$  holds.

Corollary 1 to Lemma 3.1 now applies to these elements  $a_1, \dots, a_m$  and this completes the proof of Lemma 3.3.

COROLLARY. *Suppose that the hypotheses (i), (ii), (iii), of Lemma 3.3 hold and suppose also that*

- (vi)  $[0, a_1 \dot{\cup} a_2]$  is upper  $\aleph$ -complete and upper  $\aleph$ -continuous.

Then  $[0, a]$  is upper  $\aleph$ -complete and upper  $\aleph$ -continuous.

*Proof.* (vi) implies (iv), (v). Hence  $[0, a]$  is upper  $\aleph$ -complete by Lemma 3.3. Upper  $\aleph$ -continuity then follows from [1, Theorem 4.3].

LEMMA 3.4. (*Additivity of lower  $\aleph$ -continuity*). *Suppose that  $[0, a_1 \cup \dots \cup a_m]$  is lower  $\aleph$ -complete and that  $[0, a_i]$  is lower  $\aleph$ -*

continuous for  $i=1, \dots, m$ . Then  $[0, a_1 \cup \dots \cup a_m]$  is lower  $\aleph$ -continuous.

*Proof.* We may assume that  $\{a_1, \dots, a_m\}$  is an independent set (replace  $a_i$  by  $[a_i - (a_i \cap (a_1 \cup \dots \cup a_{i-1}))]$  for  $2 \leq i \leq m$ ).

Then  $[a_1, a_1 \cup a_2]$  is lower  $\aleph$ -continuous since it is lattice isomorphic to  $[0, a_2]$  under the mapping:  $x \rightarrow x \cap a_2$ . Similarly  $[a_2, a_1 \cup a_2]$  is lower  $\aleph$ -continuous. By the dual of [1, Theorem 4.3],  $[0, a_1 \cup a_2] = ([a_1 \cap a_2, a_1 \cup a_2])$  is lower  $\aleph$ -continuous. Lemma 3.4 follows by induction on  $m$ .

**LEMMA 3.5.** *Suppose that each of  $[0, a \cup b]$ ,  $[0, b \cup c]$ ,  $[0, a \cup c]$  is lower  $\aleph$ -complete and suppose that  $l(a, c, \aleph)$  holds. Then  $[0, a \cup b \cup c]$  is lower  $\aleph$ -complete.*

*Proof.* We may suppose that  $\{a, b, c\}$  is an independent set, for if  $c, b$  are replaced by  $[c - (a \cap c)]$  and  $[b - (b \cap (a \cup c))]$  respectively the hypotheses of Lemma 3.5 continue to hold ( $l(a, c, \aleph)$  is equivalent to  $l(a, c, \aleph)$  if  $a \cup c_1 = a \cup c$ ) and the conclusion is not changed.

Now set  $B = a \cup c$ ,  $C = b \cup a$ ,  $A = b \cup c$ , and  $1 = a \cup b \cup c$ . We have:  $[A \cap B, 1] (= [c, a \cup b \cup c])$  is lower  $\aleph$ -complete since it is lattice isomorphic to  $[0, a \cup b]$  under the mapping  $x \rightarrow x \cap (a \cup b)$ . Similarly each of  $[B \cap C, 1]$ ,  $[C \cap A, 1]$  is lower  $\aleph$ -complete.

We can now show that  $[0, a \cup b \cup c] (= [A \cap B \cap C, 1])$  is lower  $\aleph$ -complete (by applying the dual of Lemma 3.1) if we can show:

(3.4) *Whenever  $X_\alpha \geq C \cap A$  for  $\alpha \in I$  (with  $\bar{I} \leq \aleph$ ) and  $C \cup (\bigcap (X_\beta | \beta \in F)) = 1$  for all finite  $F \subset I$ , then  $C \cup (\bigcap (X_\alpha | \alpha \in I)) = 1$ .*

Since  $C \cap A = b$  and  $C = a \cup b$ , (3.4) can be rewritten:

(3.4)' *Whenever  $X_\alpha \geq b$  for  $\alpha \in I$  (with  $\bar{I} \leq \aleph$ ) and  $a \cup (\bigcap (X_\beta | \beta \in F)) = a \cup b \cup c$  for all finite  $F \subset I$  then  $a \cup (\bigcap (X_\alpha | \alpha \in I)) = a \cup b \cup c$ .*

Suppose that the hypotheses of (3.4)' hold and set  $x_\alpha = X_\alpha \cap (a \cup c)$ . Then  $x_\alpha \leq a \cup c$  for all  $\alpha$  and

$$\begin{aligned} & a \cup (\bigcap (x_\beta | \beta \in F)) \\ &= a \cup ((\bigcap (X_\beta | \beta \in F)) \cap (a \cup c)) = (a \cup (\bigcap (X_\beta | \beta \in F))) \cap (a \cup c) \\ &= (a \cup b \cup c) \cap (a \cup c) = a \cup c. \end{aligned}$$

Since  $l(a, c, \aleph)$  holds, it follows that

$$\begin{aligned} & a \cup (\bigcap (x_\alpha | \alpha \in I)) = a \cup c; \quad a \cup (\bigcap (X_\alpha | \alpha \in I) \cap (a \cup c)) = a \cup c; \\ & a \cup (\bigcap (X_\alpha | \alpha \in I)) \geq a \cup c \text{ (hence } = a \cup b \cup c \text{)}. \end{aligned}$$

This means: (3.4)' does hold. This completes the proof of Lemma 3.5.

COROLLARY 1. Suppose that  $[0, a_i \cup a_j]$  is lower  $\mathfrak{K}$ -complete for  $i, j = 1, \dots, m$ .

Suppose also that  $l(a_i, a_j, \mathfrak{K})$  holds for all  $i < j$ . Then  $[0, a_1 \cup \dots \cup a_m]$  is lower  $\mathfrak{K}$ -complete.

*Proof.* This follows from Lemma 3.5 by induction on  $m$ , just as Corollary 1 to Lemma 3.1 followed from Lemma 3.1.

COROLLARY 2. Suppose that  $[0, a_i \cup a_j]$  is lower  $\mathfrak{K}$ -complete and lower  $\mathfrak{K}$ -continuous for  $i, j = 1, \dots, m$ . Then  $[0, a_1 \cup \dots \cup a_m]$  is lower  $\mathfrak{K}$ -continuous.

*Proof.* Since lower  $\mathfrak{K}$ -continuity of  $[0, a_i \cup a_j]$  implies that  $l(a_i, a_j, \mathfrak{K})$  holds, Corollary 1 shows that  $[0, a_1 \cup \dots \cup a_m]$  is lower  $\mathfrak{K}$ -complete. The lower  $\mathfrak{K}$ -continuity of  $[0, a_1 \cup \dots \cup a_m]$  then follows from Lemma 3.4.

LEMMA 3.6. Suppose that

- (i)  $a = a_1 \cup a_2 \cup \dots \cup a_m$  for some finite  $m \geq 2$ ,
- (ii)  $a_2 \sim a_1$ ,
- (iii)  $a_i \lesssim a_1 \cup \dots \cup a_{i-1}$  for  $2 < i \leq m$ ,
- (iv)  $[0, a_1 \cup a_2]$  is lower  $\mathfrak{K}$ -complete,
- (v)  $l(a_1, a_2, \mathfrak{K})$  holds.

Then  $[0, a]$  is lower  $\mathfrak{K}$ -complete.

COROLLARY. Suppose that (i), (ii), (iii) hold and also

- (vi)  $[0, a_1 \cup a_2]$  is lower  $\mathfrak{K}$ -complete and lower  $\mathfrak{K}$ -continuous.

Then  $[0, a]$  is lower  $\mathfrak{K}$ -complete and lower  $\mathfrak{K}$ -continuous.

*Proof.* Lemma 3.6 and its Corollary follow from Lemma 3.5 and Lemma 3.4 just as Lemma 3.3 and its Corollary followed from Corollary 1 to Lemma 3.1 and [1, Theorem 4.3].

THEOREM 3.1. Suppose that each of  $[0, a_i \cup a_j]$  is an  $\mathfrak{K}$ -von Neumann-geometry (respectively a von Neumann-geometry) for  $i, j = 1, \dots, m$ . Then  $[0, a_1 \cup \dots \cup a_m]$  is an  $\mathfrak{K}$ -von Neumann-geometry (respectively a von Neumann geometry).

*Proof.* This follows from Corollary 2 to Lemma 3.1 and Corollary 2 to Lemma 3.5.

COROLLARY 1. Suppose that

- (i)  $a = a_1 \cup a_2 \cup \dots \cup a_m$  for some finite  $m \geq 2$ ,

- (ii)  $a_2 \sim a_1$ ,
- (iii)  $a_i \lesssim a_1 \dot{\cup} \dots \dot{\cup} a_{i-1}$  for  $2 < i \leq m$ ,
- (iv)  $[0, a_1 \dot{\cup} a_2]$  is an  $\aleph$ -von Neumann-geometry (respectively a von Neumann-geometry).

Then  $[0, a]$  is an  $\aleph$ -von Neumann-geometry, respectively a von Neumann-geometry.

*Proof.* This follows from the Corollary to Lemma 3.3 and the Corollary to Lemma 3.6.

**COROLLARY 2.** Suppose that  $\mathcal{R}$  is an  $\aleph$ -von Neumann-ring (respectively a von Neumann-ring). If  $\bar{R}_{\mathcal{R}}$  has a basis  $x_1, x_2, \dots, x_m$  such that  $x_2 \sim x_1$  and  $x_i \lesssim x_1$  for  $2 < i \leq m$ , then  $\mathcal{R}_2$  is an  $\aleph$ -von Neumann-ring (respectively, a von Neumann-ring).

*Proof.* By hypothesis, the unit element of the lattice  $\bar{R}_{\mathcal{R}}$  is the union  $x_1 \dot{\cup} \dots \dot{\cup} x_m$ . The unit element of  $\bar{R}_{\mathcal{S}}$ , with  $\mathcal{S} = \mathcal{R}_2$ , can be represented as a union  $x_1 \dot{\cup} \dots \dot{\cup} x_m \dot{\cup} y_1 \dot{\cup} \dots \dot{\cup} y_m$  with  $y_i \sim x_i$  and hence  $y_i \lesssim x_1$  for  $1 \leq i \leq m$ . Since  $[0, x_1 \dot{\cup} x_2]$  is an  $\aleph$ -von Neumann geometry (respectively a von Neumann geometry) along with  $\bar{R}_{\mathcal{R}}$ , Corollary 1 applies and this completes the proof of Corollary 2.

**COROLLARY 3.** Suppose that  $\mathcal{R}$  and  $\mathcal{R}_2$  are both  $\aleph$ -von Neumann-rings (respectively von Neumann-rings). Then  $\mathcal{R}_n$  is an  $\aleph$ -von Neumann-ring (respectively a von Neumann-ring) for all finite  $n$ .

*Proof.* If  $n > 2$  the unit element of  $\bar{R}_{\mathcal{S}}$ , with  $\mathcal{S} = \mathcal{R}_n$ , can be expressed as  $x_1 \dot{\cup} x_2 \dot{\cup} \dots \dot{\cup} x_n$  where  $x_1$  is the unit element of  $\bar{R}_{\mathcal{R}}$ ,  $x_i \sim x_1$  for all  $i$ , and  $[0, x_1 \dot{\cup} x_2] = \bar{R}_{\mathcal{R}_2}$ . Theorem 3.1 applies and this completes the proof of Corollary 3.

**REMARK.** Let  $\mathcal{R}$  be the ring of sequences  $x = (x^n)$  with all  $x^n$  complex numbers and all but a finite number of  $x^n$  real, with componentwise addition and multiplication; this example was given by Kaplansky [3, page 526]. This  $\mathcal{R}$  is a von Neumann-ring but  $\mathcal{R}_2$  is not even upper  $\aleph_0$ -complete.

**DEFINITION 3.1.** If  $L$  is a relatively complemented modular lattice, then an element  $a$  is called Boolean (with respect to  $L$ ) if  $b_1 \sim b_2$ ,  $b_1 \leq a$  together imply  $b_1 = b_2$ ;  $a$  is called the Boolean part of  $L$  (necessarily unique if it exists)<sup>2</sup> if  $a$  is Boolean and  $a_1 \leq a$  for every Boolean  $a_1$ .

<sup>2</sup> This is an abuse of language: properly,  $[0, a]$  should be called the Boolean part of  $L$ .



LEMMA 3.7. *Suppose that  $L$  is a relatively complemented modular lattice. If  $(a, b)P$  holds then for every  $c$  in  $L$ ,  $c \cap (a \cup b) = (c \cap a) \cup (c \cap b)$  and  $[0, a \cup b]$  is the direct sum of  $[0, a]$  and  $[0, b]$ . On the other hand if  $a$  is Boolean then*

- (i)  $b \leq a$  implies that  $b$  is Boolean,
- (ii)  $b \cap a = 0$  implies that  $(b, a)P$  holds,
- (iii)  $b \geq a$  implies that the relative complement  $[b - a]$  is unique,
- (iv)  $a \cap (b \cup c) = (a \cap b) \cup (a \cap c)$  for all  $b, c$  in  $L$ ,
- (v)  $[0, a]$  is a Boolean algebra.

*Proof.* Suppose that  $(a, b)P$  holds and set  $d = [(c \cap (a \cup b)) - ((c \cap a) \cup (c \cap b))]$ ,  $d_a = (d \cup b) \cap a$ ,  $d_b = (d \cup a) \cap b$ . Then  $d \leq a \cup b$ ,  $d \cap a = d \cap b = 0$ ,  $d_a \dot{\cup} d = (d \cup b) \cap (d \cup a) = d_b \dot{\cup} d$ , so  $d_a \sim d_b$ . Since  $d_a \leq a$ ,  $d_b \leq b$  and  $(a, b)P$  holds, we must have:  $d_a = 0$ ;  $b = d_a \cup b = d \cup b$ ;  $d \leq b$ ; hence  $d = 0$ ,  $c \cap (a \cup b) = (c \cap a) \cup (c \cap b)$ . If  $c \leq a \cup b$  then  $c = (c \cap a) \cup (c \cap b)$ ; and if  $c = c_1 \cup c_2$  with  $c_1 \leq a$ ,  $c_2 \leq b$  then  $c \cap a = c_1 \cup (c_2 \cap b \cap a) = c_1 \cup 0 = c_1$ ,  $c \cap b = c_2$ . This proves that  $[0, a \cup b]$  is the direct sum of  $[0, a]$  and  $[0, b]$ .

(i) and (ii) are obvious from the definition of Boolean element.

(ii) asserts that  $a$  is in the centre of  $L$  as defined in [1, (2.5)]. But if  $a$  is in the centre of  $L$  and  $b$  is any element in  $L$  with  $b \geq a$  then  $a$  is in the centre of  $[0, b]$ , hence  $[b - a]$  is uniquely determined (use [1, (2.6)]). This proves (iii).

If  $b, c$  are arbitrary elements in  $L$ , set  $b_1 = [b - (a \cap b)]$ ,  $c_1 = [c - (a \cap c)]$ . Since  $a \cap b_1 = a \cap c_1 = 0$  and  $a$  is in the centre of  $L$ , it follows that  $(a, b_1)P$ ,  $(a, c_1)P$ , hence  $(a, b_1 \cup c_1)P$  (use [1, (2.6)]); therefore  $a \cap (b_1 \cup c_1) = 0$ . By the modular law

$$\begin{aligned} a \cap (b \cup c) &= a \cap (b_1 \cup c_1 \cup (a \cap b) \cup (a \cap c)) \\ &= (a \cap b) \cup (a \cap c) \cup (a \cap (b_1 \cup c_1)) \\ &= (a \cap b) \cup (a \cap c) \end{aligned}$$

and hence (iv) holds.

Thus  $[0, a]$  is a distributive complemented lattice, equivalently: a Boolean algebra. This proves (v).

LEMMA 3.8. *Suppose that  $L$  has a unit element  $1 = a_1 \cup a_2 \cup \dots \cup a_m$  with  $m \geq 2$ ,  $a_2 \sim a_1$ ,  $a_i \lesssim a_1$  for  $2 < i \leq m$  and  $a_1 \cap a_2 = 0$ . Then the Boolean part of  $L$  exists and is 0.*

*Proof.* By Lemma 3.2 we may assume that  $1 = a_1 \dot{\cup} \dots \dot{\cup} a_m$  with  $m \geq 2$ ,  $a_2 \sim a_1$  and  $a_i \lesssim a_1$  for  $2 < i \leq m$ .

To prove Lemma 3.8 we may suppose that  $a \neq 0$  and we need only exhibit elements  $b_1, b_2$  such that  $b_1 \leq a$ ,  $b_1 \sim b_2$ , and  $b_1 \neq b_2$ .

If  $a_i \cap a \neq 0$  for any  $i$  it suffices to choose this element as  $b_1$  since the relations  $a_1 \sim a_2$  and  $a_i \lesssim a_1$  if  $i \neq 1$  imply  $b_1 \sim b_2$  for some  $b_2 \neq b_1$  (even  $b_1 \cap b_2 = 0$ ).

On the other hand, if  $a_i \cap a = 0$  for all  $i$ , set  $b_1 = (a_1 \cup \dots \cup a_i) \cap a$  where  $i$  is the smallest integer for which this element is different from 0 (necessarily  $1 < i \leq m$ ) and set  $b_2 = ((a_1 \cup \dots \cup a_{i-1}) \cup b_1) \cap a_i$ . Then  $b_1 \sim b_2$  since  $(a_1 \cup \dots \cup a_{i-1}) \cup b_1 = (a_1 \cup \dots \cup a_{i-1}) \cup b_2$ ; and  $b_1 \neq b_2$  since  $b_2 \leq a_i$  and  $b_1 \cap a_i \leq a \cap a_i = 0$ . This completes the proof of Lemma 3.8.

**LEMMA 3.9.** *Suppose that  $L$  is an upper complete complemented modular lattice and let  $a$  be the union of all Boolean elements in  $L$ . Then  $a$  is the Boolean part of  $L$ .*

*Proof.* We need only show that  $a$  is Boolean, that is, we may suppose that  $b \leq a$ , that  $\varphi$  is a perspective mapping of  $[0, b]$ , that  $b \neq \varphi(b)$  and we need only derive a contradiction. By replacing  $b$  by  $[b - (b \cap \varphi(b))]$  we may suppose  $b \neq 0$  and  $b \cap \varphi(b) = 0$ .

Now for every  $c$ :  $(\varphi(b \cap c)) \sim (b \cap c)$  and  $(\varphi(b \cap c)) \cap (b \cap c) = 0$ . If  $c$  is Boolean this implies:  $b \cap c = 0$ , and hence (since  $c$  is Boolean)  $(b, c)P$  holds. It follows from [1, formula (2.6)] that  $(b, a)P$  holds, contradicting the fact that  $b \neq 0$  and  $b \leq a$ . This contradiction proves Lemma 3.9.

**THEOREM 3.2.** *Suppose that  $L$  is a relatively complemented modular lattice and*

- (i)  $a = a_0 \cup a_1 \cup a_2 \cup \dots \cup a_m$  for some finite  $m \geq 2$ ,
- (ii)  $(a_0, a_1 \cup \dots \cup a_m)P$  holds,
- (iii)  $a_2 \sim a_1, a_2 \cap a_1 = 0$ ,
- (iv)  $a_i \lesssim a_1 \cup \dots \cup a_{i-1}$  for  $2 < i \leq m$ ,
- (v)  $\varphi$  is a perspective mapping of  $[0, b]$  with  $\varphi(b) \leq a$ .

Let  $\pi$  denote one of the properties: to be upper  $\aleph$ -complete and upper  $\aleph$ -continuous, or to be lower  $\aleph$ -complete and lower  $\aleph$ -continuous. Then  $[0, a \cup b]$  has property  $\pi$  if both of  $[0, a_1 \cup a_2]$  and  $[0, a_0 \cup \varphi^{-1}(a_0 \cap \varphi(b))]$  have property  $\pi$ ; if  $a_0$  is the Boolean part of  $[0, a]$  and  $[0, b]$  has a Boolean part  $b_0$ , it is sufficient that  $[0, a_1 \cup a_2]$  and  $[0, a_0 \cup b_0]$  should both have property  $\pi$ .

*Proof.* Since  $(a_0, a_1 \cup \dots \cup a_m)P$  holds, Lemma 3.7 shows that  $\varphi(b) = \varphi(b_1) \cup \varphi(b_2)$  where  $b_1 = \varphi^{-1}(a_0 \cap \varphi(b))$  and  $b_2 = \varphi^{-1}((a_1 \cup \dots \cup a_m) \cap \varphi(b))$ . Then  $(a_0 \cup b_1, a_1 \cup \dots \cup a_m \cup b_2)P$  holds (use [1, (2.6)]).

By Lemma 3.7,  $[0, a \cup b]$  is the direct sum of  $[0, a_0 \cup b_1]$  and  $[0, a_1 \cup \dots \cup a_m \cup b_2]$  and has property  $\pi$  if each of the summands has it.

Since  $b_2 \lesssim a_1 \cup \dots \cup a_m$ ,  $[0, a_1 \cup \dots \cup a_m \cup b_2]$  has property  $\pi$  if  $[0, a_1 \cup a_2]$  has it, by Lemma 3.3 and its Corollary and Lemma 3.6 and its Corollary.

If  $a_0$  is the Boolean part of  $[0, a]$  then  $\varphi(b) \cap a_0$  is Boolean with respect to  $[0, a]$ , a fortiori Boolean with respect to  $[0, \varphi(b)]$ . Thus,  $b_1$  is Boolean with respect to  $[0, b]$ . If  $[0, b]$  has a Boolean part  $b_0$  then  $b_1 \leq b_0$  and  $a_0 \cup b_1 \leq a_0 \cup b_0$ , hence  $[0, a_0 \cup b_1]$  has property  $\pi$  if  $[0, a_0 \cup b_0]$  has it.

This proves all parts of Theorem 3.2.

REMARK. If  $\mathcal{R}$  is a von Neumann ring then  $\mathcal{R}$  has a unique decomposition as a direct sum  $\mathcal{R} = \mathcal{B} \oplus \mathcal{R}$  such that  $\bar{R}_{\mathcal{R}}$  is the Boolean part of  $\bar{R}_{\mathcal{R}}$  and  $\bar{R}_{\mathcal{R}}$  has a basis  $x_1, x_2, x_3$  with  $x_2 \sim x_1$  and  $x_3 \lesssim x_1$ . Then Theorem 3.2 and Corollary 2 to Theorem 3.1 apply and show that  $\mathcal{R}_2$  is a von Neumann ring if and only if  $\mathcal{B}_2$  is a von Neumann ring (for details see [2]).

#### REFERENCES

1. Ichiro Amemiya and Israel Halperin, *Complemented modular lattices*, Canadian J. of Math., **11** (1959), 481-520.
2. Israel Halperin, *Elementary divisors in von Neumann rings*, Acta Scientiarum Mathematicarum Szeged, to appear.
3. Irving Kaplansky, *Any orthocomplemented complete modular lattice is a continuous geometry*, Annals of Math., **61** (1955), 524-541.
4. John von Neumann, *Continuous Geometry*, Princeton University Press, 1960.

QUEEN'S UNIVERSITY,  
KINGSTON, CANADA.

